

**GLOBAL OCEAN SURFACE WATER PARTIAL PRESSURE OF CO<sub>2</sub> DATABASE:  
MEASUREMENTS PERFORMED DURING 1957–2010  
(Version 2010)**

Contributed by  
Taro Takahashi and Stewart C. Sutherland

Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY 10964

Date Published: September 2007  
Date Updated: June 2011

Prepared for the Climate Change Research Division  
Office of Biological and Environmental Research  
U.S. Department of Energy  
Budget Activity Numbers KP 12 04 01 0 and KP 12 02 03 0

Prepared by  
Alex Kozyr

Carbon Dioxide Information Analysis Center  
OAK RIDGE NATIONAL LABORATORY  
Oak Ridge, Tennessee 37831-6335  
managed by  
UT-BATTELLE, LLC  
for the  
U.S. DEPARTMENT OF ENERGY  
under contract DE-AC05-00OR22725



## CONTENTS

List of Figures.....	v
List of Tables .....	v
Abbreviations and Acronyms .....	vii
Abstract.....	ix
1. INTRODUCTION .....	1
2. OBJECTIVE.....	2
3. SUMMARY OF ADDITIONS AND CORRECTIONS IN VERSION 2010 .....	2
4. METHODS OF COMPUTATION .....	3
5. QUALITY CONTROL.....	5
6. DATA SOURCES.....	6
7. DATA LISTING.....	7
8. HOW TO OBTAIN THE DATA AND DOCUMENTATION .....	9
9. ACKNOWLEDGEMENTS.....	9
10. REFERENCES .....	9



## **LIST OF FIGURES**

1	Location of LDEO V2010 master database of sea surface pCO <sub>2</sub> observations. ....	2
---	---	---

## **LIST OF TABLES**

1	List of data contributors to the global surface water pCO <sub>2</sub> LDEO database .....	6
2	Data elements listed in the master data file “LDEO_databaseV2010.txt” .....	8
3	Data elements listed in the metadata file “LDEO_Database_Metadata_V2010.xls” .....	8



## ABBREVIATIONS AND ACRONYMS

AOML	Atlantic Oceanographic and Meteorological Laboratory
CDIAC	Carbon Dioxide Information Analysis Center
CLIVAR	Climate Variability (Program)
CO <sub>2</sub>	carbon dioxide
CSIRO	Australian Commonwealth Scientific and Research Organization
DOE	U.S. Department of Energy
FTP	file transfer protocol
$f$ CO <sub>2</sub>	fugacity of CO <sub>2</sub>
GEOSECS	Geochemical Ocean Sections Study
IGY	International Geophysical Year
JGOFS	Joint Global Ocean Flux Study
LDEO	Lamont-Doherty Earth Observatory
NCAR	National Center for Atmospheric Research
NCEP	National Centers for Environmental Prediction
NDP	numeric data package
NOAA	National Oceanic and Atmospheric Administration
NSF	National Science Foundation
ORNL	Oak Ridge National Laboratory
pCO <sub>2</sub>	partial pressure of CO <sub>2</sub>
PMEL	Pacific Marine Environmental Laboratory
SSS	sea surface salinity
SST	sea surface temperature
VOS	Volunteer Observing Ship
WOCE	World Ocean Circulation Experiment





## ABSTRACT

**Takahashi, T., S.C. Sutherland, and A. Kozyr. 2011. Global Ocean Surface Water Partial Pressure of CO<sub>2</sub> Database: Measurements Performed During 1957–2010 (Version 2010). ORNL/CDIAC-159, NDP-088(V2010). Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee, doi: 10.3334/CDIAC/otg.ndp088(V2010).**

Approximately 5.2 million measurements of surface water partial pressure of CO<sub>2</sub> obtained over the global oceans during 1957–2010 are listed in the Lamont-Doherty Earth Observatory (LDEO) database, which includes open ocean and coastal water measurements. The data assembled include only those measured by equilibrator-CO<sub>2</sub> analyzer systems and have been quality-controlled based on the stability of the system performance, the reliability of calibrations for CO<sub>2</sub> analysis, and the internal consistency of data. To allow re-examination of the data in the future, a number of measured parameters relevant to pCO<sub>2</sub> measurements are listed. The overall uncertainty for the pCO<sub>2</sub> values listed is estimated to be  $\pm 2.5$   $\mu\text{atm}$  on the average.

For simplicity and for ease of reference, this version is referred to as V2010, meaning that data collected through 31 December 2010 has been included. It is our intention to update this database annually. In this update, 21 new cruise/ship files are added to the previous version V2009.

The data presented in this database include the analyses of partial pressure of CO<sub>2</sub> (pCO<sub>2</sub>), sea surface temperature (SST), sea surface salinity (SSS), pressure and temperature of the equilibration, and barometric pressure in the outside air from the ship's observation system. A minor equilibration temperature problem has been corrected in the R/V *Laurence M. Gould* data for 2010.

The global pCO<sub>2</sub> data set is available free of charge as a numeric data package (NDP) from the Carbon Dioxide Information Analysis Center (CDIAC). The NDP consists of the oceanographic data files and this printed documentation, which describes the procedures and methods used to obtain the data.

**Keywords:** carbon dioxide, partial pressure of CO<sub>2</sub>, global carbon cycle, global ocean, equilibrator-CO<sub>2</sub> analyzer systems.




## 1. INTRODUCTION

Transfer of CO<sub>2</sub> from the atmosphere to the oceans is a critical process in the global carbon cycle, and is important for the future of the earth's climate. During the past decade, about 6 Pg-C yr<sup>-1</sup> (1 Pg (peta grams) = 10<sup>15</sup> grams = 1 Giga ton) has been emitted into the atmosphere from various anthropogenic sources including the combustion of fossil fuels, cement production, deforestation, land use changes and others. On the basis of various independent scientific methods including changes in oxygen and CO<sub>2</sub> concentrations in the atmosphere (Keeling et al. 1996; Bender et al., 2005; Manning and Keeling, 2006), sea-air differences in CO<sub>2</sub> partial pressure (Takahashi et al., 2002; Takahashi et al., 2009), distribution of carbon isotopes (<sup>12</sup>C, <sup>13</sup>C and <sup>14</sup>C) (Gruber and Keeling, 2001; Quay et al, 2003), inversion of atmospheric CO<sub>2</sub> distribution data using atmospheric circulation models (Baker et al., 2006; Patra et al., 2006), and various global carbon cycle models (Sarmiento et al., 2000; Gruber and Sarmiento, 2002; Matsumoto et al., 2004; Gruber et al., 2009), the annual uptake rate of CO<sub>2</sub> by the oceans has been estimated to be about 2 Pg-C yr<sup>-1</sup> for the past decade. Thus, about 30% of the anthropogenic CO<sub>2</sub> emissions are absorbed annually by the oceans, and about 50% remain in the atmosphere. As a result, the atmospheric CO<sub>2</sub> concentration is increasing at a mean rate of about 1.5 ppm yr<sup>-1</sup> (or 0.4% per year), and the concentration of CO<sub>2</sub> dissolved in surface ocean waters is also increasing, thus causing the acidification of ocean waters. Accurate documentation of changes that have occurred in the CO<sub>2</sub> chemistry of ocean waters is therefore important for understanding the fate of anthropogenic CO<sub>2</sub> released into the atmosphere as well as charting the future course of atmospheric CO<sub>2</sub> levels that would affect the earth's climate.

Carbon dioxide molecules react chemically with water to form bicarbonate (HCO<sub>3</sub><sup>-</sup>) and carbonate (CO<sub>3</sub><sup>=</sup>) ions, both of which do not communicate with the overlying air. Only about 0.5% of the total CO<sub>2</sub> molecules in seawater communicate with the air via gas exchange across the sea surface. This quantity is called the partial pressure of CO<sub>2</sub> (pCO<sub>2</sub>), which represents the CO<sub>2</sub> vapor pressure. The seawater CO<sub>2</sub> depends on the temperature, the total amount of CO<sub>2</sub> dissolved in seawater and the pH of seawater. Over the global ocean, it varies from about 100 μatm to 1000 μatm (1 μatm = 10<sup>-6</sup> atm = 0.101325 Pascal). When seawater pCO<sub>2</sub> is smaller than the atmospheric pCO<sub>2</sub> (presently about 390 μatm), seawater takes up CO<sub>2</sub> from the overlying air; when it is greater than the atmospheric pCO<sub>2</sub>, it emits CO<sub>2</sub> to the air. The rate of transfer of CO<sub>2</sub> across the sea surface is estimated by: (sea-air CO<sub>2</sub> flux) = (transfer coefficient) x (sea-air p CO<sub>2</sub> difference). The transfer coefficient depends primarily on the degree of turbulence near the interface, and is commonly expressed as a function of the square of wind speed. Since the time-space variation for atmospheric pCO<sub>2</sub> is much smaller than that for surface ocean water pCO<sub>2</sub>, the magnitude of sea-air pCO<sub>2</sub> difference and hence the net flux is governed primarily by seawater pCO<sub>2</sub>. Therefore, the interannual and seasonal variability for surface water pCO<sub>2</sub> is of particular interest.

The atmospheric CO<sub>2</sub> concentrations observed at a number of locations over the globe is summarized in GLOBALVIEW- CO2 (2010) and TRENDS ON LINE (2010). On the other hand, no single data file for surface water pCO<sub>2</sub> that includes long-term and global coverage has been made accessible to the general public. About 30 years ago, we started to assemble a global surface water pCO<sub>2</sub> data for time-space variability studies for the global oceans using the observations made by the Lamont-Doherty Earth Observatory (LDEO) group. Many investigators from other institutions contributed data to the database for the first publication on the global ocean pCO<sub>2</sub> and sea-air CO<sub>2</sub> flux (Takahashi et al. 1997), which was based on about 0.25 million pCO<sub>2</sub> measurements; the second publication (Takahashi et al. 2002) was based on about a million measurements. In the third publication in the Deep Sea Research (Takahashi et al., 2009), a dataset that consisted of about 3 million pCO<sub>2</sub> measurements plus a number of other measured supporting parameters, has been analyzed and summarized. A subset of this database is used for estimating the sea-air CO<sub>2</sub> flux from coastal waters surrounding North America (Chavez and Takahashi, 2007). The LDEO database V2010 now consists of more than 5.2 million pCO<sub>2</sub> measurements (Fig. 1) plus a number of other measured supporting parameters. Mindful of the increasing

importance of CO<sub>2</sub> studies for future global welfare, LDEO is sharing this data file widely with the global research communities and the public through the Carbon Dioxide Information Analysis Center (CDIAC) data archive.



**Fig. 1. Location of LDEO V2010 master database of sea surface pCO<sub>2</sub> observations.**

## **2. OBJECTIVE**

Modern investigation of ocean water pCO<sub>2</sub> started during the International Geophysical Year (IGY), 1957–1960, with the introduction of infrared CO<sub>2</sub> gas analyzer and gas-water equilibrator aboard oceanographic research vessels (Takahashi, 1961; Keeling et al., 1965; Keeling, 1965; Broecker and Takahashi, 1966; Keeling and Waterman, 1968; Keeling, 1968; Waterman et al., 2006-a, b, c and d). Combined with high-accuracy CO<sub>2</sub>-air-gas mixtures made available for the calibrations of the analyzers at sea, the quality of seawater pCO<sub>2</sub> data was greatly improved from that obtained in the pre-World War II era. During major oceanographic programs that took place after IGY, the CO<sub>2</sub> analyzers and equilibrators have been steadily improved to give more frequent observations and better accuracy, although the principles of the measurement remained essentially unchanged. Computers used for system control and data logging contributed significantly to the data quality. The objective of this database is to assemble high-quality pCO<sub>2</sub> data obtained using the equilibrator-analyzer method, and to reprocess the data using a standardized method of computation, that will be described below. Therefore, the pCO<sub>2</sub> values listed in this database may differ from the original listings prepared by respective investigators.

## **3. SUMMARY OF ADDITIONS AND CORRECTIONS IN VERSION 2010**

This version is referred to as Version 2010, and includes the data collected through 31 December 2010. In this update, 21 new cruise/ship files are added to the previous Version 2009 (see NDP-088\_V2009).

Other than routine updates from our Vessel of Opportunity Program such as *L.M. Gould*, *N.B. Palmer*, and *R/V Ronald H. Brown*, there are two new sources of data: *R/V Marcus G. Langseth*, operated by Lamont-Doherty Earth Observatory, and *S.A. Agulhas* operated by CSIR, South Africa.

#### 4. METHODS OF COMPUTATION

The seawater pCO<sub>2</sub> data listed in LDEO database are based on direct measurements of seawater pCO<sub>2</sub> made using equilibrator-CO<sub>2</sub> analyzer systems. Many of the data listed are from semi-continuous pCO<sub>2</sub> systems with flow-through water; many others are measurements for discrete water samples made at hydrographic stations along with measurements for other chemical and physical properties. Although different types of equilibrators (e. g., shower type, bubbling type, membrane type, and rotating disks in flow-through or closed circulation systems) and CO<sub>2</sub> gas analyzers (non-dispersive infrared analyzers and gas chromatographs of various designs) were employed, the results from different systems are accepted as long as analyzers were properly calibrated using validated CO<sub>2</sub>-air gas mixtures and the carrier gas was equilibrated with seawater samples. Because of the diversity of methods used, it is impossible to present details of the method used by each contributing research group. Detailed methodology may be obtained directly from the investigators listed in Table 1, or from the CDIAC reports for specific expeditions.

It is important to point out that the methods used for computing CO<sub>2</sub> concentrations in equilibrated gas have been varied among groups. For example, some groups computed a least-square fit of readings for three or more standard gas mixtures to a quadratic equation and used it to calculate concentrations in samples. Some groups used four or five standard gas mixtures for calibrations, and fitted the data to a 4<sup>th</sup> order polynomial equation. And some groups used an output from linearization circuits of infrared analyzer, and linearly regressed three or more standard gas readings to obtain sample CO<sub>2</sub> concentrations. The outputs from a gas chromatograph are a linear function of CO<sub>2</sub> concentration, and hence a linear regression was used for calibration. These different data reduction methods yield CO<sub>2</sub> concentrations varying  $\pm 1.5$  ppm (or  $\pm 1.5$   $\mu$ atm in pCO<sub>2</sub>). However, we did not recompute the CO<sub>2</sub> values using a single uniform algorithm but instead, accepted CO<sub>2</sub> concentration values as reported to us. Since different analyzers and different numbers of standard gas mixtures were used by respective groups, no single uniform data reduction scheme can be applied, and hence we relied on the judgment of each group for selecting the data reduction scheme most suited for their operational modes and skills. Measurements that were made using only one calibration gas mixture were judged unreliable and were not included in this database.

Using the reported CO<sub>2</sub> concentration values, the pCO<sub>2</sub> value in sample seawater at the equilibration temperature, (pCO<sub>2</sub>)<sub>eq</sub>, has been recomputed with the relationship:

$$(pCO_2)_{eq} = V_{co2} (P_{eq} - P_{water}),$$

where  $V_{co2}$  is the mole fraction concentration of CO<sub>2</sub> in carrier gas ( $V_{CO2}$  is same as  $X_{CO2}$ , which is often used in literature, and these qualities may be used interchangeably);  $P_{eq}$  is the total pressure of gas in the equilibrator; and  $P_{water}$  is the equilibrium water vapor pressure at temperature of equilibration,  $T_{eq}$ , and salinity. Since some equilibrators were operated open to the room air,  $P_{eq}$  values may be equal to the ship's interior pressure or to the barometric pressure outside the ship depending on the location of the equilibrator. When an equilibrator is located in an enclosed shipboard laboratory and is open to the room air,  $P_{eq}$  is the ambient pressure in the laboratory. While an equilibrator operated in an enclosed space, only the barometric pressure at sea surface was reported in some data sets, but not  $P_{eq}$ . In such cases,  $P_{eq}$  is assumed to be the reported barometric pressure at sea surface plus 3 mb, that represents an overpressure normally maintained inside a ship. This correction increases the (pCO<sub>2</sub>)<sub>sw</sub> value by about 1  $\mu$ atm. When the pressure was not reported, we used the climatological value in the nearest box from the National Centers for Environmental Prediction/National Center for Atmospheric Research (NCEP/NCAR)

Reanalysis II Project file for the month of the observation. The  $p\text{CO}_2$  at in-situ seawater temperature is computed using an integrated form of the temperature effect for isochemical seawater,  $(\partial \ln p\text{CO}_2 / \partial T)_{\text{Sal, Alk, TCO}_2}$  (Takahashi et al., 1993):

$$(p\text{CO}_2)_{\text{sw}} @ T_{\text{in situ}} = [(p\text{CO}_2)_{\text{sw}} @ T_{\text{eq}}] \text{Exp}\{0.0433 (T_{\text{in situ}} - T_{\text{eq}}) - 4.35 \times 10^{-5} [(T_{\text{in situ}})^2 - (T_{\text{eq}})^2]\}$$

where the “sw” and “eq” indicate the in situ and equilibrator conditions respectively. Throughout the computation,  $\text{CO}_2$  gas is assumed to behave as an ideal gas that mixes with air and water vapor ideally. Although  $\text{CO}_2$  fugacity is used in a number of published papers and data reports, we refrained from using the fugacity since it is computed differently from an investigator to another. Although we do not list the sea-air  $p\text{CO}_2$  differences in this report, we recommend the formula below for the computation of atmospheric  $p\text{CO}_2$  and the corresponding value for sea-air  $p\text{CO}_2$  difference.

$$(p\text{CO}_2)_{\text{air}} = (V_{\text{CO}_2})_{\text{air}} (P_{\text{baro}} - P_{\text{sw}})$$

where  $P_{\text{baro}}$  is the barometric pressure at sea surface, and  $P_{\text{sw}}$  is the equilibrium water vapor pressure at the temperature and salinity for mixed layer water. The subscript “air” indicates the value for atmosphere samples.

The sea-air  $p\text{CO}_2$  difference,  $\Delta p\text{CO}_2$ , is then computed using:

$$\Delta p\text{CO}_2 = (p\text{CO}_2)_{\text{sw}} - (p\text{CO}_2)_{\text{air}}$$

Since  $\text{CO}_2$  is assumed to be an ideal gas for both  $(p\text{CO}_2)_{\text{sw}}$  and  $(p\text{CO}_2)_{\text{air}}$ , the small effects of non-ideality should cancel due to differencing for  $\Delta p\text{CO}_2$ . Positive  $\Delta p\text{CO}_2$  values indicate that the sea is a source for atmospheric  $\text{CO}_2$ , whereas negative values indicate that the sea is a sink.

Beginning with V2007 we added a column reporting the partial pressure of  $\text{CO}_2$  in seawater in units of Pascals.

Values for the fugacity of  $\text{CO}_2$  in seawater,  $f\text{CO}_2$ , have been submitted to LDEO by some investigators. In principle, the chemical potential of  $\text{CO}_2$  should be represented by  $f\text{CO}_2$ . However, in practice, the fugacity values reported by various investigators are not clearly defined as to whether only the non-ideality arising from  $\text{CO}_2$ - $\text{CO}_2$  molecular interactions is considered and/or that from  $\text{CO}_2$ - $\text{H}_2\text{O}$  interactions is also included. The original determination of the solubility of  $\text{CO}_2$  in seawater by Murray and Riley (1971), which was conducted by equilibrating 1 atmosphere total pressure of pure  $\text{CO}_2$  gas, appears to include implicitly the effects of  $\text{CO}_2$ - $\text{H}_2\text{O}$  interactions in gas phase above seawater samples, since the  $\text{CO}_2$  gas above the sample water contained water vapor in equilibrium with the sample water at respective temperatures. Weiss (1974) refined their data and presented a formulation for the  $\text{CO}_2$  solubility under 1 atmosphere  $\text{CO}_2$  fugacity. However, while Weiss's results account primarily for the  $\text{CO}_2$ - $\text{CO}_2$  interactions, it is probable that the effect of  $\text{CO}_2$ - $\text{H}_2\text{O}$  interactions in the gaseous phase is also included. Subsequently, Weiss and Price (1980) proposed a formulation for the  $\text{CO}_2$ - $\text{H}_2\text{O}$  interactions based upon Pitzer's correlations for interactions among gas molecules. Some researchers (e. g. Wanninkhof and Thoning, 1993) chose to correct Weiss's fugacity values using, in addition, the Weiss-Price  $\text{CO}_2$ - $\text{H}_2\text{O}$  non-ideality effect. Whether such a procedure may constitute a double correction for the  $\text{CO}_2$ - $\text{H}_2\text{O}$  interactions is not clear. Because of these ambiguities, we have chosen to report  $p\text{CO}_2$  values assuming ideality. Since the mole fraction concentrations of  $\text{CO}_2$  in equilibrated gas samples, temperature and pressure of equilibration are also included in the file, anyone who wishes to compute  $f\text{CO}_2$  will be able to compute it with one's preferred algorithm for fugacity.

Since  $f\text{CO}_2$  values are always smaller than the corresponding  $p\text{CO}_2$  values by 1 to 2  $\mu\text{atm}$  and the differences are large enough with respect to the precision of measurements and the mean global sea-air  $p\text{CO}_2$  difference of about 10  $\mu\text{atm}$ , they should not be used interchangeably with  $p\text{CO}_2$ . On the other hand, the sea-air difference in  $f\text{CO}_2$  is numerically indistinguishable with measurement errors from the

sea-air pCO<sub>2</sub> difference (pCO<sub>2</sub>) as long as fCO<sub>2</sub> in air and pCO<sub>2</sub> in air are computed in the same manner as fCO<sub>2</sub> and pCO<sub>2</sub> for the seawater respectively.

In all OISO data files, only fCO<sub>2</sub> values are reported without pCO<sub>2</sub> values or CO<sub>2</sub> concentrations in dried equilibrated gas. We had to convert the reported fCO<sub>2</sub> values to pCO<sub>2</sub> values to make the OISO data consistent with the rest of the database. Since formulations used for fCO<sub>2</sub> calculation are not provided in their data report, it is not possible to re-compute pCO<sub>2</sub>. Therefore, we use the following approximation for the conversion;

$$pCO_2 (\mu atm) = fCO_2 (\mu atm) \times [1.00436 - 4.669 \times 10^{-5} \times SST (^{\circ}C)]$$

This yields pCO<sub>2</sub> values greater than fCO<sub>2</sub> by 0.8 to 1.8  $\mu atm$  (about 1.3  $\mu atm$  on the average) in the range of temperatures and pCO<sub>2</sub> values encountered in the OISO study areas in the southern Indian Ocean.

## 5. QUALITY CONTROL

Determination of surface water pCO<sub>2</sub> requires measurements for a number of parameters in addition to the concentration of CO<sub>2</sub> in the carrier gas equilibrated with seawater. With the broad range of data sources and continued evolution of the measurement systems, it has not been possible to ensure that the observations contain all the necessary data elements with desired precisions. In some cases, pCO<sub>2</sub> or fCO<sub>2</sub> was reported without V<sub>CO<sub>2</sub></sub>, sometimes with the pressure at the time of equilibration, sometimes not. Some files contained only the sea surface temperature and the pCO<sub>2</sub> at that temperature, but not the temperature and pCO<sub>2</sub> at equilibration. Ideally, the incoming file should contain the dried concentration of CO<sub>2</sub> in a parcel of air in equilibrium with seawater (V<sub>CO<sub>2</sub></sub>), the temperature at the time of equilibration, and the equilibration pressure. When the pressure was not reported, we used the climatological value in the nearest box from the NCEP/NCAR Reanalysis II Project file for the month of the observation. We accepted data points that contained pCO<sub>2</sub> and sea surface temperature with or without the pressure.

Field data obtained in earlier years do not necessarily have all the required supporting measurements. For example, the temperatures of equilibration and water samples were recorded by hand at intervals of a few to several hours, and hence these values are often interpolated linearly with time between measurements. The salinity was measured only at hydrographic stations, and these values were interpolated between stations. When salinity was not measured during some expeditions, climatological values were assigned. Some data were unrecoverable and lost as a result of changes in computer systems (hardware and software). For the past decade, computer controlled systems for underway surface water pCO<sub>2</sub> measurements have been deployed widely, and more complete high quality data sets have been obtained. In constructing this data file, we exercised our personal judgment for accepting and rejecting data, especially those collected in earlier years. For more recent data sets, water flow rates through the equilibrator and the temperatures of equilibrator water are recorded for each pCO<sub>2</sub> measurement. They often served as important criteria for identifying satisfactory operation of the equilibrator. Measurements made at reduced or stopped water flow conditions were rejected, as were those made at unusually rapid changes in the temperature of the water in an equilibrator. CO<sub>2</sub> gas analyzers are commonly calibrated using three or more gas mixtures of different CO<sub>2</sub> concentrations. Those measurements made using only a single calibration gas mixture are subject to unspecified uncertainties, and hence are rejected.

All data points have been examined individually as they were integrated into this database. While not perfect, we have tried to edit out obvious problems, erring on the side of leaving in questionable ones. Considering differences in equilibrator designs, calibration methods and some interpolated parameters, we estimate that the uncertainty of the pCO<sub>2</sub> data presented in this data file is about  $\pm 2.5 \mu atm$  on the average.

Seawater pCO<sub>2</sub> values that were computed using the alkalinity, total CO<sub>2</sub> concentration, and pH data are not included in this data file because of their large uncertainties and potential biases resulting from different dissociation constants of carbonic, boric and other acids used for the computations.

During many cruises, atmospheric CO<sub>2</sub> concentrations were measured concurrently with surface water pCO<sub>2</sub>. However, we suspect that many of these measurements were contaminated by local sources (such as ship's exhausts). Since atmospheric CO<sub>2</sub> concentrations vary from one air mass to another, especially in the northern hemisphere, distinguishing local contamination from natural variability is difficult. Hence, we decided to omit the atmospheric CO<sub>2</sub> data from this file. We recommend that the atmospheric CO<sub>2</sub> concentration data listed in the GLOBALVIEW-CO2 for computing the sea-air pCO<sub>2</sub> difference.

## 6. DATA SOURCES

The LDEO database lists more than 5.2 million surface ocean pCO<sub>2</sub> observations made since 1957. A large portion of the data is composed of observations from the continuous underway systems, such as those used aboard the National Science Foundation icebreakers *Nathaniel B. Palmer* and *Laurence B. Gould* operating primarily in the Southern Ocean, and the research ships operated by the Atlantic Oceanographic and Meteorological Laboratory and the Pacific Marine Environmental Laboratory of the National Oceanic and Atmospheric Administration. The database also includes the observations made during a number of major national and international oceanographic programs such as Geochemical Ocean Sections Study (GEOSECS), Joint Global Ocean Flux Study (JGOFS), World Ocean Circulation Experiment (WOCE), Climate Variability (CLIVAR) Repeat Hydrography Project, Volunteer Observing Ships (VOS) Project and others, which were supported by the National Science Foundation, National Oceanic and Atmospheric Administration, and Department of Energy. A significant number of data have been contributed by international colleagues from many countries including Japan, Germany, France, UK, Iceland, Australia, Canada, the Netherlands, Norway, and others. Table 1 lists the major contributors for the database.

**Table 1. List of data contributors to the global surface water pCO<sub>2</sub> LDEO database.**

PI name <sup>*</sup>	Institution	Country
Thorarinn S. Arnarson	Marine Research Institute and University of Iceland	Iceland
Dorothee C. E. Bakker	School of Environmental Sciences, University of East Anglia	UK
Nicholas R. Bates	Bermuda Institute of Ocean Sciences	Bermuda
Richard Bellarby	Bjerknes Centre for Climate Research, University of Bergen	Norway
Wei-Jun Cai	University of Georgia	USA
Francisco Chavez	Monterey Bay Aquarium Research Institute	USA
David W. Chipman <sup>**</sup>	Lamont-Doherty Earth Observatory, Columbia University	USA
Cathy E. Cosca	Pacific Marine Environmental Laboratory, National Oceanographic and Atmospheric Administration	USA
Bruno Delille	Universite de Liege, Liege	Belgium
Hein J. W. de Baar	Netherland Institute for Sea Research	Netherlands
Richard A. Feely	Pacific Marine Environmental Laboratory, National Oceanographic and Atmospheric Administration	USA
Gernot Friederich	Monterey Bay Aquarium Research Institute	USA
John Goddard <sup>**</sup>	Lamont-Doherty Earth Observatory, Columbia University	USA
Burke Hales	College of Oceanic and Atmospheric Sciences, Oregon State University	USA
Mario Hoppema	Alfred Wegener Institute for Polar and Marine Research	Germany
Masao Ishii	Meteorological Research Institute	Japan
Truls Johannessen	Bjerknes Centre for Climate Research, University of Bergen	Norway
C. D. Keeling <sup>***</sup>	Scripps Institution of Oceanography, University of California San Diego	USA



Arne Körtzinger	Leibniz Institute of Marine Sciences	Germany
Nicolas Metzl	Laboratoire d’Oceanographie et du Climat, Universite Pierre et Marie Curie	France
Takashi Midorikawa	Meteorological Research Institute	Japan
Ludger Mintrop	MARIANDA marine analytics and data	Germany
Pedro M. S. Monteiro	CSIR, South Africa, Jan Cilliers St, Stellenbosch 7599	South Africa
P. P. Murphy	Pacific Marine Environmental Laboratory, National Oceanographic and Atmospheric Administration	USA
Timothy Newberger	Lamont-Doherty Earth Observatory, Columbia University	USA
Yukihiro Nojiri	National Institute for Environmental Studies	Japan
Jon Olafsson	Marine Research Institute and University of Iceland	Iceland
Are Olsen	Bjerknes Centre for Climate Research, University of Bergen	Norway
Christopher L. Sabine	Pacific Marine Environmental Laboratory, National Oceanographic and Atmospheric Administration	USA
Ute Schuster	School of Environmental Sciences, University of East Anglia, Norwich	UK
Tobias Steinhoff	Leibniz Institute of Marine Sciences	Germany
Stewart C. Sutherland	Lamont-Doherty Earth Observatory, Columbia University	USA
Peter Salomeh	Scripps Institution of Oceanography	USA
Colm Sweeney	Earth System Research Laboratory, National Oceanographic and Atmospheric Administration	USA
Taro Takahashi	Lamont-Doherty Earth Observatory, Columbia University	USA
Bronte Tilbrook	Australian Commonwealth Scientific and Research Organization (CSIRO) Marine and Atmospheric Research	Australia
Rik Wanninkhof	Atlantic Oceanographic and Meteorological Laboratory, National Oceanographic and Atmospheric Administration	USA
Lee S. Waterman**	Scripps Institution of Oceanography, University of California San Diego	USA
Andrew Watson	School of Environmental Sciences, University of East Anglia	UK
Ray F. Weiss	Scripps Institution of Oceanography, University of California San Diego	USA
C. S. Wong	Institute of Ocean Sciences	Canada
H. Yoshikawa-Inoue	Graduate School of Environmental Earth Science, Hokkaido University	Japan

\*The PI names are given in the alphabetical order.

\*\*Retired

\*\*\*Deceased

## 7. DATA LISTING

This NDP consists of two files: the file called “LDEO\_Database\_V2010.txt” contains all the numerical data, and the file “LDEO\_Database\_Metadata\_V2010.csv” contains information on the data source, credit, institution, etc. The data elements in “LDEO\_Database\_V2010.txt” and their units are listed in Table 2; and those in “LDEO\_Database\_Metadata\_V2010.csv” are explained in Table 3. Detailed cruise documentation (such as names of ship and ports and dates for departure and arrival) may be obtained directly from the respective PI’s.

**Table 2. Data elements listed in the master data file “LDEO\_Database\_V2009.txt”**

CRUISE_ID*	Internal LDEO Cruise Number
STN*	Station number assigned in the LDEO File
LAT	Latitude in decimal degrees (North is positive)
LON	Longitude in decimal degrees (East is positive)
MONTH/DAY/YEAR	Date
JDATE	Julian Date in decimal notation. Convention is 0001 UTC 1 Jan = 1.0
VCO2_SW**	Mole fraction concentration of CO <sub>2</sub> (ppm) in dried air
TEMP_PCO2	Temperature at which pCO <sub>2</sub> was measured in °C
SST	Sea Surface Temperature in °C
SSS	Sea Surface Salinity
PCO2_SST	Partial Pressure of CO <sub>2</sub> in seawater (in units of microatmospheres) at the temperature in the SST column
PCO2_SSTPA	Partial Pressure of CO <sub>2</sub> in seawater (in units of Pascals) at the temperature in the TEMP column
PCO2_TEQ	Partial Pressure of CO <sub>2</sub> in seawater (in units of microatmospheres) at the temperature in the TEMP_PCO2 column. This is ordinarily the value that is actually measured
EQ_PBARO	Pressure in the equilibration vessel in units of millibars
SHIPBARO	Barometric pressure in the outside air from the ship's observation system in units of millibars

\*Cross reference fields to the internal LDEO file name in case problems are discovered

\*\*Missing data have the value -999.9

**Table 3. Data elements listed in the metadata file “LDEO\_Database\_Metadata\_V2009.csv”**

CruiseID	Internal LDEO File Name
LEG	Leg Number. This is an arbitrary designation in many files because of our limit of four characters for a station number. For surface underway files that normally have many more than 9,999 observations we have broken the file into “legs.”
SHIP/EXPERIMENT	The name of the ship or other platform used, and experiment, project
CRUISE_NAME	Cruise name
OBSERVER	A notation of the person making the observation or responsible for running the collection system
DEPARTURE_PORT	Port of Departure
DEPARTURE_DATE	Date of Departure
ARRIV_EPORT	Port of Arrival
ARRIV_EDATE	Date of Arrival
COMMENT*	Air data source and general information

\*Additional information may be available for a particular file if needed.

## 8. HOW TO OBTAIN THE DATA AND DOCUMENTATION

The LDEO database NDP-088(V2010) is available free of charge from CDIAC. The complete documentation and data can be obtained from the CDIAC oceanographic web site ([http://cdiac.ornl.gov/oceans/LDEO\\_Underway\\_Database/](http://cdiac.ornl.gov/oceans/LDEO_Underway_Database/)).

Contact information:

Carbon Dioxide Information Analysis Center  
Oak Ridge National Laboratory  
P.O. Box 2008  
Oak Ridge, Tennessee 37831-6335  
USA  
Telephone: (865) 574-3645  
Telefax: (865) 574-2232  
E-mail: [cdiac@ornl.gov](mailto:cdiac@ornl.gov)  
Internet: <http://cdiac.ornl.gov/>

## 9. ACKNOWLEDGEMENTS

Many national and international investigators contributed their observations to this database. ORNL gratefully acknowledges the efforts of investigators and supporting staff at sea and on land. Richard A. Feely [Pacific Marine Environmental Laboratory/National Oceanographic and Atmospheric Administration (PMEL/NOAA)] and Rik Wanninkhof [Atlantic Oceanographic and Meteorological Laboratory (AOML) /NOAA] offered encouragement in this task; this was appreciated. . We gratefully acknowledge that the continued support from the Office of Global Programs of the National Oceanic and Atmospheric Administration has made this compilation possible. The field study of T. Takahashi and his associates have been supported by the National Science Foundation, the National Oceanic and Atmospheric Administration, Department of Energy, Comer Science and Education Foundation, Ford Motor Company and EXXON Research and Engineering Company.

## 10. REFERENCES

- Baker D. F., R. M. Law, K. R. Gurney, P. Rayner, P. Peylin, A. S. Denning, P. Bousquet, L. Bruhwiler, Y.-H. Chen, P. Ciais, I. Y. Fung, M. Heimann, J. John, T. Maki, S. Maksyutov, K. Masari, M. Prather, B. Pak, S. Taguchi, and Z. Zhu. 2006. TransCom 3 inversion intercomparison: Impact of transport model errors on the interannual variability of regional CO<sub>2</sub> fluxes, 1988-2003. *Global Biogeochem. Cycles*, 20, GB1002, doi. 10.1029/2004GB2002439.
- Bender, M.L., D.T. Ho, M.B. Hendricks, R. Mika, M.O. Battle, P.P. Tans T.J. Conway, B. Sturtevant, and N. Cassar. 2005. "Atmospheric O<sub>2</sub>/N<sub>2</sub> changes, 1993–2002: Implications for the partitioning of fossil fuel CO<sub>2</sub> sequestration," *Global Biogeochemical Cycles* 19:GB4017, doi:10.1029/2004GB002410.

- Broecker, W.S. and T. Takahashi. 1966. „Calcium carbonate precipitation on the Bahama Banks,“ *Journal of Geophysical Research* 71:1575–1602.
- Chavez, F. and T. Takahashi. 2007. Coastal oceans, Chapter 15, in “The First State of the Carbon Cycle Report (SOCCR): North American Carbon Budget and Implications for the Global Carbon Cycle”. A report by the U.S. Climate Change Science Program and the Subcommittee on Global Change Research [King, A.W. L. Dilling, G.P. Zimmerman, D.M. Fairman, R.A. Houghton, G.H. Marland, A.Z. Rose, and T.J. Wilbanks (eds.)] National Ocean and Atmospheric Administration, Climate Program Office, Silver Spring, MD, USA, pp. 83–92.
- GLOBALVIEW-CO2: 2006. *Cooperative Atmospheric Data Integration Project—Carbon Dioxide*. CD-ROM, NOAA CMDL, Boulder, Colorado (also available on Internet via anonymous FTP to ftp.cmdl.noaa.gov, Path: ccg/co2/GLOBALVIEW).
- Gruber, N., and C. D. Keeling. 2001. An improved estimate of the isotopic air-sea disequilibrium of CO<sub>2</sub>: Implications for the oceanic uptake of anthropogenic CO<sub>2</sub>. *Geophys. Res. Lett.*, 28(3), 555–558, doi:10.1029/2000GL011853.
- Gruber, N. and J.L. Sarmiento. 2002. “Large-scale biogeochemical-physical interactions in elemental cycles.” In *The Sea*, Vol. 12:337–99. John Wiley & Sons, New York.
- Gruber, N., M. Gloor, S. E. Mikaloff Fletcher, S. C. Doney, S. Dutkiewicz, M. Follows, M. Gerber, A. R. Jacobson, F. Joos, K. Lindsay, D. Menemenlis, A. Mouchet, S. A. Müller, J. L. Sarmiento, and T. Takahashi. 2009. Oceanic sources, sinks, and transport of atmospheric CO<sub>2</sub>. *Glob. Biogeochem. Cycles.*, 23, GB1005, doi:10.1029/2008GB003349.
- Keeling, C. D. 1965. Carbon dioxide in surface waters of the Pacific Ocean, 2, Calculation of the exchange with the atmosphere. *Journal of Geophysical Research*, 70:6099–102.
- Keeling, C. D., N. W. Rakestraw and L. S. Waterman. 1965. Carbon dioxide in surface waters of the Pacific Ocean, 1, Measurements of the distribution. *Journal of Geophysical Research*, 70:6087–98.
- Keeling, C. D. 1968. Carbon dioxide in surface waters, 4. Global distribution. *Journal of Geophysical Research*, 73:4529–4542.
- Keeling, C. D. and L.S. Waterman. 1968. Carbon dioxide in surface waters, 3. Measurements on Lusiad Expedition 1962-1963. *Journal of Geophysical Research*, 73:4529–42.
- Keeling, R.F., S.C. Piper, and M. Heinmann. 1996. “Global and hemispheric CO<sub>2</sub> sinks deduced from changes in atmospheric O<sub>2</sub> concentration,” *Nature* 381:218–21.
- Manning, A. C. and R. F. Keeling. 2006. Global oceanic and land biotic carbon sinks from the Scripps atmospheric oxygen flask sampling network. *Tellus B* 58:95–116.
- Murray, C. N. and J. P. Riley. 1971. The solubility of gases in distilled water and seawater - IV. Carbon dioxide. *Deep-Sea Res.*, 18:533–541.

- Matsumoto, K. et al. 2004. Evaluation of ocean carbon cycle models with data-based metrics. *Geophys. Res. Lett.*, 31, LO7303, doi:10.1029/2003GL018970.
- Metzl, N. 2009. Decadal increase of oceanic carbon dioxide in the Southern Indian Ocean surface waters (1991–2007). *Deep-Sea Res. II*, doi:10.1016/j.dsr2.2008.12.007.
- Patra, P. K., Gurney, K. R., Denning, A. S., Maksyutov, S., Nakazawa, T., Baker, D., Bousquet, P., Bruhwiler, L., Chen, Y.-H., Ciais, P., Fan, S., Fung, I., Gloor, M., Heinmann, M., Higuchi, K., John, J., Law, R. M., Maki, T., Pak, B.C., Peylin, P., Prather, M., Rayner, P. J., Sarmiento, J., Tauchi, S., Takahashi, T. and Yuen, C.- W. 2006. Sensitivity of inverse estimation of annual mean CO<sub>2</sub> sources and sinks to ocean-only sites versus all-sites observational networks. *Geophys. Res. Lett.*, 33, LO5814, doi:10.1029/2005GL025403.
- Quay, P., R. Sommerup, T. Westby, J. Sutsman, and A. McNichol. 2003. “Changes in the <sup>13</sup>C/<sup>12</sup>C of dissolved inorganic carbon in the ocean as a tracer of anthropogenic CO<sub>2</sub> uptake.” *Global Biogeochemical Cycles* 17, No. 1:DOI 10.1029/2001GB001817.
- Sarmiento, J. L., P. Monfray, E. Maier-Reimer, O. Aumont, R.J. Murnane, and J.C. Orr. 2000. “Sea-air CO<sub>2</sub> fluxes and carbon transport: A comparison of three ocean general circulation models.” *Global Biogeochemical Cycles* 14:1267–81.
- Sabine, C.L., R. A. Feely, N. Gruber, R. M. Key, K. Lee, J. L. Bullister, R. Wanninkhof, C. S. Wong, D. W. R. Wallace, B. Tilbrook, F. J. Millero, T.-H. Peng, A. Kozyr, T. Ono, A. F. Rios. 2004. The Oceanic Sink for Anthropogenic CO<sub>2</sub>. *Science* 305(5682):367-371.
- Takahashi, T. 1961. “Carbon dioxide in the atmosphere and in the Atlantic Ocean water,” *Journal of Geophysical Research* 66:477–94.
- Takahashi, T., J. Olafsson, J. Goddard, D.W. Chipman, and S.C. Sutherland. 1993. “Seasonal variation of CO<sub>2</sub> and nutrients in the high-latitude surface oceans: A comparative study,” *Global Biogeochemical Cycles* 7:843–78.
- Takahashi, T., R.A. Feely, R. Weiss, R. Wanninkhof, D.W. Chipman, S.C. Sutherland, and T. Takahashi. 1997. “Global air-sea flux of CO<sub>2</sub>: an estimate based on measurements of sea-air pCO<sub>2</sub> difference,” *Proc. Nat. Acad. Sci.*, 94:8292–99.
- Takahashi, T., S.C. Sutherland, C. Sweeney, A. Poisson, N. Metzl, B. Tillbrook, N. Bates, R. Wanninkhof, R.A. Feely, C.L. Sabine, J. Olafsson, and Y. Nojiri. 2002. “Global sea-air CO<sub>2</sub> flux based on climatological surface ocean pCO<sub>2</sub>, and seasonal biological and temperature effects,” *Deep-Sea Research II*, 49:1601–22.
- Takahashi, T., S. C. Sutherland, R. Wanninkhof, C. Sweeney, R. A., Feely, D. W. Chipman, B. Hales, G. Friederich, F. Chavez, C. Sabine, A. Watson, D. C. E. Bakker, U. Schuster, N. Metzl, H. Yoshikawa-Inoue, M. Ishii, T. Midorikawa, Y. Nojiri, A. Kortzinger, T. Steinhoff, M. Hoppema, J. Olafsson, T. S. Arnarson, B. Tillbrook, T. Johannessen, A. Olsen, R. Bellerby, C. S. Wong, B. Delille, N. R. Bates, and H. J. W. de Baar. 2009. Climatological mean and decadal change in surface ocean pCO<sub>2</sub> and net sea-air CO<sub>2</sub> flux over the global oceans. *Deep-Sea Res. II*, 56, 554–577, doi:10.1016/j.dsr2.2008.12.009.

- TRENDS on line. 2006. "A Compendium of Data on Global Change. Carbon Dioxide Information and Analysis Center," Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, TN <http://cdiac.ornl.gov/trends/co2/contents/html>).
- Wanninkhof, R. and K.Thoning. 1993. Measurement of fugacity of CO<sub>2</sub> in surface water using continuous and discrete sampling methods. *Marine Chem.*, 44:189–204.
- Waterman, L. S., P. P. Tans, T. Aten, C. D. Keeling, and T. A. Boden. 2006-a. Surface water and atmospheric underway carbon dioxide data obtained during Downwind (1957) expedition of Scripps Institution of Oceanography. Scripps Institution of Oceanography, Univ. California San Diego, La Jolla, CA  
[http://scrippsco2.ucsd.edu/data/seawater\\_pco2/pco2\\_downwind\\_1957.csv](http://scrippsco2.ucsd.edu/data/seawater_pco2/pco2_downwind_1957.csv)
- Waterman, L. S., P. P. Tans, T. Aten, C. D. Keeling, and T. A. Boden. 2006-b. Surface water and atmospheric underway carbon dioxide data obtained during " Monsoon (1961) expedition of Scripps Institution of Oceanography. Scripps Institution of Oceanography, Univ. California San Diego, La Jolla, CA  
[http://scrippsco2.ucsd.edu/data/seawater\\_pco2/pco2\\_monsoon\\_1961.csv](http://scrippsco2.ucsd.edu/data/seawater_pco2/pco2_monsoon_1961.csv)
- Waterman, L. S., P. P. Tans, T. Aten, C. D. Keeling, and T. A. Boden. 2006-c. Surface water and atmospheric underway carbon dioxide data obtained during Lusiad (1962) expedition of Scripps Institution of Oceanography. Scripps Institution of Oceanography, Univ. California San Diego, La Jolla, CA  
[http://scrippsco2.ucsd.edu/data/seawater\\_pco2/pco2\\_lusiad\\_1962.csv](http://scrippsco2.ucsd.edu/data/seawater_pco2/pco2_lusiad_1962.csv)
- Waterman, L. S., P. P. Tans, T. Aten, C. D. Keeling, and T. A. Boden. 2006-d. Surface water and atmospheric underway carbon dioxide data obtained during Lusiad (1963) expedition of Scripps Institution of Oceanography. Scripps Institution of Oceanography, Univ. California San Diego, La Jolla, CA  
[http://scrippsco2.ucsd.edu/data/seawater\\_pco2/pco2\\_lusiad\\_1963.csv](http://scrippsco2.ucsd.edu/data/seawater_pco2/pco2_lusiad_1963.csv)
- Weiss, R. F. 1974. Carbon dioxide in water and seawater: The solubility of a non-ideal gas. *Marine Chem.*, 2:203–215.
- Weiss, R. F. and B. A. Price. 1980. Nitrous oxide solubility in water and seawater. *Marine Chem.*, 8:347–359.